



Southwest General

Partnering with




University Hospitals

EMS Services

PRE-HOSPITAL CARE

MEDICAL CONTROL

PROTOCOLS AND PROCEDURES



ACUTE CORONARY SYNDROME

**UNIVERSAL PATIENT CARE
PROTOCOL**

B	EMT-B	B
I	EMT-I	I
P	EMT-P	P
M	MED CONTROL	M

Oxygen
10-15 L
Titrate to > 94% SpO₂

Apply Cardiac Monitor

Go to Appropriate Dysrhythmia

Obtain 12 – Lead EKG
(Look for ST Elevation and transmit
to the hospital)

CHEST PAIN AND EKG INDICATES STEMI

Use caution with acute inferior wall MI
(II, III, AVF) – place IV prior to Nitroglycerine. Normal
Saline bolus prior to Nitroglycerine strongly recommended
Use caution with acute septal wall MI
(V1,V2) – watch for AV blocks–consider placing pacing pads

IV Protocol

ASPIRIN
324 mg chew and swallow
(81 mg / tab x 4)

Hypotension / Dysrhythmias
Treat per Appropriate Protocol

NITROGLYCERIN 0.4 mg SL every 5 minutes x 3
(If BP greater than 90 Systolic with IV)
(If BP greater than 110 Systolic without IV)
*Basic EMT's may assist pt. with 1 of their
own nitro.

(if no relief with a total of 3 NTG)
Morphine Sulfate
2 mg IV every 4-5 minutes titrated to
respiratory status and pain (MAX = 10 mg)

Reassess and Monitor

Continued Pain?

Consider Nitrous Oxide if no relief from
Morphine

CONTACT MEDICAL CONTROL

ACUTE CORONARY SYNDROME

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Age • Medications • Past medical history (MI, Angina, Diabetes) • Recent physical exertion • Onset Palpatations 	<ul style="list-style-type: none"> • CP (pain, pressure, aching, vice like tightness) • Location (substernal, arm, jaw, epigastric, neck, shoulder) • Radiation of pain • Pale, diaphoresis • Shortness of breath • Nausea, vomiting, dizziness 	<ul style="list-style-type: none"> • Trauma vs. Medical • Angina vs. Myocardial infarction • Pericarditis • Pulmonary embolism • Asthma / COPD • Pneumothorax • Aortic dissection or aneurysm • GE reflux or Hiatal hernia • Esophageal spasm • Chest wall injury or pain • Pleural pain

GENERAL CONSIDERATIONS:

- Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro
- Be suspicious of a “Silent MI” in the elderly, diabetics, and women. Diabetics and geriatric patients often have atypical pain, or only generalized complaints.
- Consider other causes of chest pain such as aortic aneurysms, pericarditis, and pulmonary embolisms.
- Oxygen administration is first, 12-Lead EKG, continuous cardiac monitoring, and an IV are indicated for patient’s who’s chest pain was relieved prior to your arrival.
- All patients complaining of chest discomfort must be administered enough O2 to maintain SPO2 of greater than 94% nasal cannula. Administer oxygen by non-rebreather or assist the patient’s ventilations as indicated.
- Aspirin is administered to achieve a therapeutic dose of 324 mg (4 chewable, 81 mg tabs), unless allergic reaction or peptic ulcer disease.
- Nitroglycerin can be administered to a patient by EMS up to 3 doses. If the patient has already taken 3 of their own prior to your arrival, document if the patient had any changes in their symptoms or a headache after taking their own Nitroglycerin. **DO NOT** administer Nitroglycerin to a patient who took an erectile dysfunction medication (Viagra, Cialas, Levitra, etc.) within the last 48 / 72 hours due to potential severe hypotension.
- If patient has taken nitroglycerin without relief, consider potency of the medication. Check and document the expiration date of the patient’s prescribed nitroglycerin.
- Nitroglycerin can be administered to a hypertensive patient complaining of chest discomfort without Medical Direction permission.
- Nitroglycerin can be administered without an IV as long as the patient takes Nitroglycerin at home and has a BP greater than 110 mmHg or BP greater than 150 mmHg if over 70 years old.
- **DO NOT** treat the PVC’s with Lidocaine or Amiodarone, if the patient is bradycardic.
- If positive ECG changes, establish a second IV while en route to the hospital.
- Monitor for hypotension after administration of nitroglycerin and morphine.
- If pain continues after O₂, ASA and Nitro, administer Morphine 2 mg IV every 4-5 minutes up to 10 mg. Titrate to response and respirations.
- If the patient becomes hypotensive from Nitroglycerin administration, place the patient in the Trendelenburg position and administer a 200 - 400 mL Normal Saline bolus.

SINUS BRADYCARDIA

B	EMT-B	B
I	EMT-I	I
P	EMT-P	P
M	MED CONTROL	M

UNIVERSAL PATIENT CARE



IV Protocol



**Apply 12 – Lead EKG
(Look for ST Elevation)
Communicate to ED**



**Hypotension, BP less than 90 Systolic,
Altered Mental status, chest pain**

No

Yes

UNIVERSAL PATIENT CARE PROTOCOL

**ATROPINE 0.5 mg IV/IO
Repeat every 3-5 minutes
If Atropine ineffective**



**Consider Sedation
VALIUM
2 – 5 mg slow IV**



**EXTERNAL TRANSCUTANEOUS
PACING**



***Consider DOPAMINE while awaiting Pacer
2 - 10 mcg/kg/min IV
Titrate to BP greater than 90 systolic**

CONTACT MEDICAL CONTROL



TRANSPORT

SINUS BRADYCARDIA

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Past medical history • Medications • Beta Blockers • Calcium channel blockers • Digitalis • Pacemaker 	<ul style="list-style-type: none"> • HR less than 60 per min. • Chest pain • Respiratory distress • Hypotension • Altered mental status • Syncope 	<ul style="list-style-type: none"> • Acute MI • Hypoxia • Hypothermia • Sinus Brady • Athletes • Head Injury (elevated ICP) or Stroke • Spinal cord lesion • Sick Sinus Syndrome • AV blocks (1st, 2nd or 3rd degree)

GENERAL CONSIDERATIONS:

- Exam: Mental Status, Neck, Heart, Lungs, Neuro
- The use of lidocaine in heart block can worsen bradycardia and lead to asystole and death.
- Pharmacological treatment of Bradycardia is based upon the presence or absence of hypotension.
- If hypotension exists, treat.
- If blood pressure is adequate, monitor only.
- Transcutaneous pacing is the treatment of choice for Type II second-degree heart blocks and third degree heart blocks.
- If the patient is **critical** and an IV is not established, initiate pacing with **Medical Direction permission**.
- If the patient converts to another rhythm, refer to the appropriate protocol and treat accordingly.
- At lower doses dopamine has a more selective effect on inotropy and heart rate; higher doses (>10 mcg) also has vasoconstrictive effects.
- May use Dopamine drip as an alternative to TCP.

NARROW – COMPLEX TACHYCARDIA

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Past medical history • Medications (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin) • Diet (caffeine, chocolate) • Drugs (nicotine, cocaine) • History of palpitations / heart racing • Syncope / near syncope 	<ul style="list-style-type: none"> • HR greater than <150 per min • QRS 0.12 sec • Dizziness, CP, SOB • Potential presenting rhythm • Sinus Tachycardia • Atrial fibrillation / flutter • Multifocal atrial tachycardia 	<ul style="list-style-type: none"> • Heart disease (WPW, Valvular) • Sick Sinus Syndrome • Myocardial infarction • Electrolyte imbalance • Exertion, pain, emotional stress • Fever • Hypoxia • Hypovolemia or anemia • Drug effect / overdose • Hyperthyroidism • Pulmonary embolus

GENERAL CONSIDERATIONS:

- Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro
- Adenosine may not be effective in identifiable atrial flutter / fibrillation, yet is not harmful.
- Continuous pulse oximetry is required for all SVT patients.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.
- If the patient converts to another rhythm, refer to the appropriate protocol and treat accordingly.
- Examples of vagal maneuvers include bearing down, coughing, or blowing into a syringe. **DO NOT** perform a carotid massage.
- If possible, the IV should be initiated in either the left or right AC.
- Consider applying the Combo patches prior to Adenosine administration.
- When administering Adenosine, raise the patient’s arm and immediately follow the bolus with 20 mL rapid bolus of normal saline.
- Record 3-Lead EKG strips during Adenosine administration.
- Perform a 12-Lead EKG prior to and after Adenosine conversion or cardioversion of SVT.
- If the patient converts into ventricular fibrillation or pulseless ventricular tachycardia, immediately DEFIBRILLATE, refer to the appropriate protocol and treat accordingly. Be sure to switch the Life Pak to PADDLES before defibrillating.
- Give a copy of the EKGs and code summaries to the receiving facility upon arrival.
- Caution giving Adenosine to patients with asthma. May cause worse bronchospasm.

Synchronized Cardioversion (mono and biphasic monitors)

If:	Sequence:			
Atrial Fibrillation	120 to 200 J	300 J	360 J	
Unstable monomorphic VT	100 to 200 J	300 J	360 J	
Other SVT Atrial Flutter	50 J	100 to 200 J	300 J	360 J
Polymorphic VT (irregular form and rate) and unstable	Treat as VF with high-energy shock (defibrillation doses)			

WIDE – COMPLEX TACHYCARDIA

UNIVERSAL PATIENT CARE PROTOCOL

B	EMT-B	B
I	EMT-I	I
P	EMT-P	P
M	MED CONTROL	M

Ventricular Fibrillation Protocol

Palpate Pulse

Yes

IV PROTOCOL

Monitor Protocol / 12 Lead EKG

Assess appropriateness for clinical condition. Heart rate typically greater than 150 bpm

Stable / Regular

Stable

Unstable

ADENOSINE 6 mg IV Push followed by 20 ml NS rapid flush (Not for atrial fibrillation)

No Response 1 – 2 minutes

ADENOSINE 12 mg IV followed by 20mL NS rapid flush

No Response 1 – 2 minutes

ADENOSINE 12 mg IV followed by 20mL NS rapid flush

If V-Tach or uncertain rhythm AMIODARONE 150 mg IV mixed in 50 mL D5W (over 10 minutes)

No Response 1 – 2 minutes

If unstable, prepare for immediate Synchronized Cardioversion

Prepare for immediate Synchronized Cardioversion 50-100 J

Consider Sedation
MIDAZOLAM (VERSED) 2-4 mg IV (2mg / 2ml)
Or
MIDAZOLAM (VERSED) 5mg Atomized IN (5mg / 1ml)
Or
DIAZEPAM (VALIUM) 2.5-5 mg slow IV
Do not confuse MIDAZOLAM (VERSED) concentrations

AMIODARONE 150 mg IV mixed in 50 mL D5W (over 10 minutes)

If Torsades de pointes – give Magnesium Sulfate 2 gm IV over 5 to 60 minutes

Repeat Synchronized Cardioversion 200, 300, 360 J

CONTACT MEDICAL CONTROL

TRANSPORT

WIDE – COMPLEX TACHYCARDIA

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Past medical history / medications, diet, drugs • Syncope / near syncope • Palpitations • Pacemaker • Allergies: lidocaine / novacaine 	<ul style="list-style-type: none"> • Ventricular tachycardia on ECG (Runs or sustained) • Conscious, rapid pulse • Chest pain, shortness of breath • Dizziness • Rate usually 150 - 180 bpm for sustained V-Tach 	<ul style="list-style-type: none"> • Artifact / Device failure • Cardiac • Endocrine / Metabolic • Drugs • Pulmonary

GENERAL CONSIDERATIONS:

- Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro
- For witnessed / monitored ventricular tachycardia, try having patient cough or deliver a precordial thump.
- Polymorphic V-Tach (Torsades de Pointes) may benefit from the administration of Magnesium Sulfate.
- If the patient converts to another rhythm, refer to the appropriate protocol and treat accordingly.
- If the patient relapses back into wide complex tachycardia / ventricular tachycardia, initiate synchronized cardioversion with the joules setting that previously cardioverted the patient.
- Record EKG strips during Amiodarone administration.
- Perform a 12- Lead EKG prior to and after Amiodarone conversion or synchronized cardioversion of wide complex tachycardia / ventricular tachycardia.
- Perform a Code Summary and attach it to the patient run report.
- Be sure to treat the patient and not the monitor.
- Magnesium Sulfate can be mixed with NS or D5W.
- Amiodarone is only compatible with D5W.

Synchronized Cardioversion (mono and biphasic monitors)

If	Sequence			
Atrial Fibrillation	120 to 200 J	300 J	360 J	
Stable monomorphic VT	100 to 200 J	300 J	360 J	
Other SVT Atrial Flutter	50 J	100 to 200 J	300 J	360 J
Polymorphic VT (irregular form and rate) and unstable	Treat as VF with high-energy shock (defibrillation doses)			

CARDIAC ARREST

UNIVERSAL PATIENT CARE PROTOCOL

B	EMT-B	B
I	EMT-I	I
P	EMT-P	P
M	MED CONTROL	M

Yes ← **Criteria for Death / DNR** → Yes

Withhold Resuscitation
Contact Medical Control

Review DNR Comfort Care Guidelines
Contact Medical Control

↓ No

CPR x 5 cycles / 2 minutes or until defibrillator is available
Defibrillate patient as soon as defibrillator or AED is available

Attach Cardiac Monitor
Defibrillate patient as soon as defibrillator or AED is available

Go to Appropriate Protocol

Deliver Shock x 1 if indicated

CPR x 5 cycles / 2 minutes

Airway Protocol

Deliver Shock x 1 if indicated

Maintain CPR / Airway

Follow AED Prompts (If applicable)

Continue CPR

IV / IO Protocol

CONTACT MEDICAL CONTROL

TRANSPORT

AT ANY TIME
Return of Spontaneous Circulation
GO TO POST RESUSCITATION CARDIAC CARE PROTOCOL

CARDIAC ARREST

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Events leading to arrest • Estimated downtime • Past medical history • Medications • Existence of terminal illness • Signs of lividity, rigor mortis • DNR or Living Will 	<ul style="list-style-type: none"> • Unresponsive • Apneic • Pulseless 	<ul style="list-style-type: none"> • Medical vs. Trauma • V. fib vs. Pulseless V. tach • Asystole • Pulseless electrical activity(PEA)

GENERAL CONSIDERATIONS:

- Exam: Mental Status
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Immediately resume chest compressions after defibrillation.
- Reassess airway frequently and with every patient move.
- Maternal Arrest - treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport.
- If the patient converts to another rhythm, refer to the appropriate protocol and treat accordingly to Return of Spontaneous Circulation Protocol.
- Attempt to obtain patient history from family members or bystanders.
 - 1) estimated down time
 - 2) medical history
 - 3) complaints prior to arrest
 - 4) bystander CPR prior to EMS arrival
 - 5) AED / CPR prior to EMS arrival
- Administer Dextrose only if the patient has a Glucose Level less than 80 with associated symptoms and is to be administered as soon a hypoglycemia is determined.
- Reassess the patient if the interventions do not produce any changes.
- If indicated, refer to the Termination of Resuscitative Efforts Protocol.

During CPR Remember: *Check pulse only at end of 2 minute cycle of CPR if an organized rhythm is present on the monitor


Push hard and fast	After an advanced airway is placed, rescuers no longer deliver “cycles” of CPR. Give continuous chest compressions without pauses for breaths at least 100 / min. Give 8 - 10 breaths / min. Check rhythm every 2 min.	<i>Search for and treat possible contributing factors:</i> Hypoxia, Hypovolemia, Hydrogen Ion (acidosis), Hypo-Hyperkalemia, Hypoglycemia, Hypothermia, Toxins, Tamponade (cardiac), Tension Pneumothorax, Thrombosis (coronary, pulmonary, or CVA), Trauma
Ensure full chest recoil		
Minimize interruptions in chest compressions		
One cycle of CPR: 30 compressions then 2 breaths 5 “cycles” = 1-2 min.		
Avoid excessive ventilation Avoid hyperventilation		
Secure airway and confirm placement	Rotate compressors every 2 min. with rhythm checks	
	Minimal interruption in chest compressions (<10 sec)	

- Medications down the endotracheal tube is to be used only if IV / EJ / IO routes can not be established

Endotracheal Guidelines - Adult and Peds

- 1) Lidocaine, Epi., Atropine, Narcan and Vasopressin can be given down the ET Tube
- 2) The optimal dose of most drugs given by ET is unknown
- 3) ET drugs deliver low blood levels. All drugs are given 2-3x’s normal dose.
- 4) Instill the drug while briefly holding compressions, follow with 5 mL (smaller with neonates) of NS flush, followed by 5 positive-pressure ventilations.

ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA)



B	EMT-B	B
I	EMT-I	I
P	EMT-P	P
M	MED CONTROL	M

ASYSTOLE / PULSELESS ELECTRICAL ACTIVITY (PEA)

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Past medical history • Medications • Events leading to arrest • End stage renal disease • Estimated downtime • Suspected hypothermia • Suspected overdose • DNR or Living Will • Tricyclics • Digitalis • Beta blockers • Calcium channel blockers 	<ul style="list-style-type: none"> • Pulseless • Apneic • No electrical activity on ECG • Cyanosis 	<ul style="list-style-type: none"> • Medical or Trauma • Hypoxia • Potassium (hypo / hyper) • Acidosis • Hypothermia • Device (lead) error • Death • Hypovolemia • Cardiac tamponade • Drug overdose (Tricyclics, Digitalis, Beta blockers, calcium channel blockers) • Massive Myocardial infarction • Tension pneumothorax • Pulmonary embolus

CONSIDER TREATABLE / REVERSIBLE CAUSES	
<ul style="list-style-type: none"> • Hypovolemia • Hypoxia • Hydrogen ion (acidosis) • Hypo-hyperkalemia • Hypoglycemia • Hypothermia 	<ul style="list-style-type: none"> • Tamponade, cardiac • Tension Pneumothorax • Thrombosis (coronary or pulmonary) • Trauma • Toxins

GENERAL CONSIDERATIONS:

- Exam: Mental Status
- Always confirm asystole in more than one lead.
- Consider each possible cause listed in the differential. Survival is based on identifying and correcting the cause!
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.
- If the patient converts to another rhythm, refer to the appropriate protocol and treat accordingly.
- Early identification and treatment of reversible causes of PEA increases the chance of a successful outcome.
- Consider volume infusion for all patients in PEA. Be alert for fluid overload.
- Treat as ventricular fibrillation if you cannot differentiate between asystole and fine ventricular fibrillation.
- Medical Direction must be contacted prior to administering antidotes for all poisonings / overdoses except for narcotic overdoses.
- Dextrose 50% should only be administered to a patient with a confirmed blood glucose level less than 80 with associated symptoms.
- Vasopressin 40 units IV / IO / ET may be given x 1 to replace first or second dose of Epi.

**VENTRICULAR FIBRILLATION (V-FIB)
PULSELESS VENTRICULAR TACHYCARDIA**

**UNIVERSAL PATIENT CARE
PROTOCOL**

B	EMT-B	B
I	EMT-I	I
P	EMT-P	P
M	MED CONTROL	M

Withhold Resuscitation and review DNR Comfort Care Guidelines

Yes

Criteria for Death / DNR

Perform CPR until defibrillator is available

Apply Cardiac Monitor Defibrillator / AED

Confirm V-Fib / Pulseless V-Tach

Defibrillate 360 J or biphasic equivalent

Resume effective CPR x 5 cycles / 2 minutes, then check pulse and rhythm

AIRWAY PROTOCOL

IV / IO Protocol

Continue effective CPR x 5 cycles / 2 minutes, then check pulse and rhythm

**EPINEPHRINE 1 mg IV/IO
1:10,000 Solution
Repeat every 3 – 5 minutes
OR**

VASOPRESSIN 40 units IV / IO. Give ONE Dose

* Meds given IV / IO when available during CPR (before and after shock)

Defibrillate 360 J or biphasic equivalent

Continue effective CPR x 5 cycles / 2 minutes, then check pulse and rhythm

Give Antiarrhythmic during CPR

Defibrillate 360 J after 5 cycles of CPR or biphasic equivalent

Continue effective CPR x 5 cycles / 2 minutes, then check pulse and rhythm

Consider Termination if Jurisdiction Authorizes

CONTACT MEDICAL CONTROL

TRANSPORT

AMIODARONE
300 mg IV
May repeat @ 150 mg IV once in 3 – 5 minutes
OR

LIDOCAINE
1-1.5 mg / kg IV
Repeat 0.5-0.75 mg / kg in 3 – 5 minutes

CONSIDER

MAGNESIUM SULFATE
1-2 g slow IV
(Torsades, ONLY)

AT ANY TIME RETURN OF SPONTANEOUS CIRCULATION (ROSC) GO TO POST RESUSCITATION CARDIAC CARE PROTOCOL

ACLS/ARRHYTHMIAS

7A

**VENTRICULAR FIBRILLATION (V – FIB)
PULSELESS VENTRICULAR TACHYCARDIA**

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Estimated down time • Past medical history • Medications • Events leading to arrest • Renal failure / dialysis • DNR or Living Will 	<ul style="list-style-type: none"> • Unresponsive, apneic, pulseless • Ventricular fibrillation or ventricular tachycardia on ECG 	<ul style="list-style-type: none"> • Asystole • Artifact / Device failure • Cardiac • Endocrine / Metabolic • Drugs • Pulmonary

GENERAL CONSIDERATIONS:

- Exam: Mental Status
- Effective CPR should be as continuous as possible with a minimum of 5 cycles or 2 minutes.
- Reassess and document at least two methods of confirming endotracheal tube placement and end tidal CO₂ frequently, after every move, and at discharge.
- Polymorphic V-Tach (Torsades de Pointes) may benefit from administration of magnesium sulfate.
- If the patient converts to another rhythm, or has a return of circulation, refer to the appropriate protocol and treat accordingly.
- If the patient converts back to ventricular fibrillation or pulseless ventricular tachycardia after being converted to ANY other rhythm, defibrillate at the previous setting used.
- Defibrillation following effective CPR is the definitive therapy for ventricular fibrillation and pulseless ventricular tachycardia.
- Vasopressin 40 units IV / IO / ET may be given x1 to replace first or second dose of Epi.
- Magnesium Sulfate should be administered early in the arrest if hypomagnesium (chronic alcoholic or malnourished patients) is suspected.
- Magnesium Sulfate can be mixed with NS or D5W.
- Amiodarone is the antiarrhythmic of choice in treating VF, Pulseless VT. Lidocaine is considered an alternative if amiodarone is not available.
- Amiodarone is only compatible with D5W.

POST – RESUSCITATION CARDIAC CARE

UNIVERSAL PATIENT CARE PROTOCOL

B	EMT-B	B
I	EMT-I	I
P	EMT-P	P
M	MED CONTROL	M

Optimize ventilation and oxygenation

- Maintain oxygen saturation $\geq 94\%$
- Consider advanced airway and waveform capnography
- Do not hyperventilate

Treat hypotension (SBP <90 mm Hg)

IV / IO Protocol

**Apply Cardiac Monitor
Defibrillator / AED
Complete 12 Lead and Transmit**

Vital Signs

Hypotension

Ventricular Ectopy

Bradycardia

Consider Fluid Bolus

**DOPAMINE
2-20 mcg / kg / min IV
Titrate to effect**

**AMIODARONE
150 mg IV
mixed in 50 ml D5W
over 10 minutes if not
previously given
OR**

**LIDOCAINE
1-1.5 mg / kg / IV**

**If rhythm converts,
LIDOCAINE DRIP
2-4 mg / minute**

**Treat per Bradycardia
Protocol**

**If arrest reoccurs, revert to
appropriate protocol**

CONTACT MEDICAL CONTROL

TRANSPORT

Reversible Causes

- ~ Hypovolemia
- ~ Hypoxia
- ~ Hydrogen ion (acidosis)
- ~ Hypo-/hyperkalemia
- ~ Hypothermia
- ~ Tension pneumothorax
- ~ Tamponade, cardiac
- ~ Toxins
- ~ Thrombosis, pulmonary
- ~ Thrombosis, coronary

POST – RESUSCITATION CARDIAC CARE

History	Signs and Symptoms	Differential Diagnosis
<ul style="list-style-type: none"> • Respiratory arrest • Cardiac arrest 	<ul style="list-style-type: none"> • Return of pulse 	<ul style="list-style-type: none"> • Continue to address specific differentials associated with the original dysrhythmia

GENERAL CONSIDERATIONS:

- Exam: Mental Status, Neck, Skin, Lungs, Heart, Abdomen, Extremities, Neuro
- Most patients immediate post resuscitation will require ventilator assistance.
- The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring.
- Appropriate post-resuscitation management can best be planned in consultation with Medical Control.
- This is the period of time between restoration of spontaneous circulation and the transfer of care at the emergency department. The focus is aimed at optimizing oxygenation and perfusion.
- Post resuscitation SVT should initially be left alone, but routinely monitor the patient. Follow Narrow Complex Tachycardia Protocol or contact Medical Control if the patient becomes hypotensive.
- If the patient is profoundly bradycardic, refer to the Sinus Bradycardia Protocol and treat accordingly.
- Adequate oxygenation is the key to a good outcome. Maintain $SPO_2 \geq 94\%$.
- Amiodarone is only compatible with D5W.
- Continuous quantitative waveform capnography is now recommended for intubated patients throughout the periarrest period. When quantitative waveform capnography is used for adults, applications now include recommendations for confirming tracheal tube placement and for monitoring CPR quality and detecting ROSC based on end-tidal carbon dioxide (PETCO₂) values
- 12 Lead completed and sent.